

Residual Inverse Iteration for the Lasing Eigenvalue Problem

Spiridonov A., Karchevskii E.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 IEEE. Effective numerical approach based on a modification of the residual inverse iteration method for solution of the Lasing Eigenvalue Problem is proposed. The numerical study demonstrates that this algorithm allows to compute all the spectral characteristics of dielectric microcavity lasers in a given frequency interval with high accuracy.

<http://dx.doi.org/10.1109/MMET.2018.8460224>

Keywords

lasing eigenvalue problem, microcavity laser, residual inverse iteration method

References

- [1] M. Lebental, E. Bogomolny, and J. Zyss, "Organic micro-lasers: A new avenue onto wave chaos physics, " in A.B. Matsko (Ed.) Practical Applications of Microresonators in Optics and Photonics, CRC Press, Boca Raton, pp. 317-353, 2009.
- [2] E. I. Smotrova and A. I. Nosich, "Mathematical study of the twodimensional lasing problem for the WG modes in a circular dielectric microcavity, " Opt. Quant. Electron, vol. 36. pp. 213-221, 2004.
- [3] E. I. Smotrova, A.I. Nosich, T.M. Benson, and P. Sewell, "Cold-cavity thresholds of microdisks with uniform and non-uniform gain: quasi-3D modeling with accurate 2D analysis, " IEEE J. Sel. Topics Quant. Electron., vol. 11, pp. 1135-1142, 2005.
- [4] E. I. Smotrova, V. Tsvirkun, I. Gozhyk, C. Lafargue, C. Ulysse, , et al., "Spectra, thresholds, and modal fields of a kite-shaped microcavity laser, " J. Opt. Soc. Am. B, vol. 30, no 6, pp. 1732-1742, 2013.
- [5] O. V. Shapoval, K. Kobayashi, and A. I. Nosich, "Electromagnetic engineering of a single-mode nanolaser on a metal plasmonic strip placed into a circular quantum wire, " IEEE J. Sel. Topics Quant. Electron., vol. 23, no 6, art no 1501609, 2017.
- [6] I. O. Sukharevsky, M. Lebental, B. Dietz, C. Lafargue, and S. Bittner, "Dielectric equilateral triangle microresonators: Integral equations and semiclassical physics approaches, " Proc. SPIE, vol. 10518, art. no 105181U-1, 2018.
- [7] A. O. Spiridonov and E. M. Karchevskii, "Field patterns of whisperinggallery and bow-Tie modes of elliptic microcavity laser with a circular active region, " Proc. Int. Conf. Transp. Opt. Networks (ICTON-2016), art. no. 7550639, 2016.
- [8] A.S. Zolotukhina, A. O. Spiridonov, E. M. Karchevskii, and A. I. Nosich "Lasing modes of a microdisk with a ring gain area and of an active microring, " Opt. Quant. Electron., vol. 47, no. 12, pp. 3883-3891, 2015.
- [9] A. S. Zolotukhina, A. O. Spiridonov, E. M. Karchevskii, and A. I. Nosich, "Electromagnetic analysis of optimal pumping of a microdisk laser with a ring electrode, " Appl. Phys. B. , vol. 123, art. no. 32, 2017.
- [10] Y.-Z. Huang, Q. Chen, W.-H. Guo, Q.-Y. Lu, and L.-J. Yu, "Mode characteristics for equilateral triangle optical resonators, " IEEE J. Sel. Top. Quant. Electron., vol. 12, no 1, pp. 59-65, 2006.

- [11] C. Lafargue, M. Lebental, A. Grigis, C. Ulysse, I. Gozhyk, N. Djellali, J. Zyss, and S. Bittner, "Localized lasing modes of triangular organic microlasers, " *Phys. Rev. E.*, vol. 90, pp. 052922/17, 2014.
- [12] J. Wiersig and M. Hentschel, "Unidirectional light emission from high-Q modes in optical microcavities, " *Phys. Rev. A*, vol. 73, no 3, pp. 031802(R)/4, 2006.
- [13] W. G. Bickley, "Two-dimension potential problems concerning a single closed boundary, " *Phil. Trans.*, vol. 228, pp 235-273, 1929.
- [14] A. O. Spiridonov, et al., "Spectra, thresholds, and modal fields of a circular microcavity laser transforming into a square, " *Proc. Int. Conf. Transp. Opt. Networks (ICTON-2015)*, art. no 7193654, 2015.
- [15] A. O. Spiridonov, et al., "Symmetry accounting in the integral-equation analysis of the lasing eigenvalue problems for two-dimensional optical microcavities, " *J. Opt. Soc. Am. B*, vol. 34, pp. 1435-1443, 2017.
- [16] A. O. Spiridonov and E. M. Karchevskii, "Mathematical and numerical analysis of the spectral characteristics of dielectric microcavities with active regions, " *Proc. Int. Conf. Days on Diffraction (DD-2016)*, pp. 390-395, 2016.
- [17] E. Karchevskii and A. Nosich, "Methods of analytical regularization in the spectral theory of open waveguides, " *Proc. Int. Conf. Math. Methods in EM Theory (MMET 2014)*, Dnipropetrovsk, pp. 39-45, 2014.
- [18] A. I. Nosich, "Method of analytical regularization in computational photonics, " *Radio Sci.*, vol. 51, no 8, pp. 1421-1430, 2016.
- [19] A. O. Spiridonov and E. M. Karchevskiy, "Projection methods for computation of spectral characteristics of weakly guiding optical waveguides, " *Proc. Int. Conf. Days on Diffraction (DD-2013)*, pp. 131-135, 2013.
- [20] A. Neumaier, "Residual inverse iteration for the nonlinear eigenvalue problem, " *SIAM J. Numer. Anal.*, vol. 22, no. 5, pp. 914-923, 1985.
- [21] I. O. Sukharevsky, A. I. Nosich, and A. Altintas, "Manipulation of backscattering from a dielectric cylinder of triangular cross-section using the interplay of GO-like ray effects and resonances, " *IEEE Trans. Antennas Propagat.*, vol. 63, no 5, pp. 2162-2169, 2015.